RESEARCH ARTICLE



WILEY

Sex-dependent clinical presentation, body image, and endocrine status in long-term remitted anorexia nervosa

Louisa Schloesser^{1,2} | Leon D. Lotter^{1,3,4,5} | Jan Offermann¹ | Katrin Borucki⁶ | Ronald Biemann⁷ | Jochen Seitz² | Kerstin Konrad^{1,8} | Beate Herpertz-Dahlmann²

¹Child Neuropsychology Section, Department of Child and Adolescent Psychiatry, Psychosomatics and Psychotherapy, University Hospital RWTH Aachen, Aachen, Germany

²Department of Child and Adolescent Psychiatry, Psychosomatics and Psychotherapy, University Hospital RWTH Aachen, Aachen, Germany

³Institute of Neuroscience and Medicine, Brain & Behaviour (INM-7), Jülich Research Centre, Jülich, Germany

⁴Institute of Systems Neuroscience, Medical Faculty, Heinrich Heine University Düsseldorf, Düsseldorf, Germany

⁵Max Planck School of Cognition, Leipzig, Germany

⁶Institute for Clinical Chemistry and Pathobiochemistry, Otto von Guericke University Magdeburg, Magdeburg, Germany

⁷Institute for Laboratory Medicine, Clinical Chemistry and Molecular Diagnostics, University Hospital Leipzig, Leipzig, Germany

⁸JARA-Brain Institute II, Molecular Neuroscience and Neuroimaging, Jülich Research Centre, Jülich, Germany

Correspondence

Louisa Schloesser, Child Neuropsychology Section, Department of Child and Adolescent Psychiatry,

Abstract

Objective: Although anorexia nervosa (AN) in males has recently gained attention, knowledge of its psychological and physiological outcomes is still scarce. We explore sex-specific characteristics of long-term remitted AN with respect to residual eating disorder (ED) psychopathology, body image, and endocrinology.

Method: We recruited 33 patients with AN in remission for at least 18 months (24 women, 9 men) and 36 matched healthy controls (HCs). Eating disorder psychopathology and body image ideals were assessed via clinical interviews, questionnaires, and an interactive 3D body morphing tool. Plasma levels of leptin, free triiodothyronine, cortisol, and sex hormones were quantified. Univariate models controlled for age and weight were used to test for the effects of diagnosis and sex.

Results: Both patient groups showed residual ED psychopathology but normal weight and hormone levels relative to HCs. Male remitted patients demonstrated significantly stronger muscularity-focused body image ideals, evident in interviews, self-reports, and behavioural data, than both female patients and HCs.

Conclusions: Sex-specific body image characteristics in patients with remitted AN point towards the need to adjust test instruments and diagnostic criteria to male-specific psychopathology. In the future, sufficiently powered studies should evaluate the risk of men with AN developing muscle dysmorphia in the long term.

KEYWORDS

anorexia nervosa, body image, hormones, remission, sex

Louisa Schloesser and Leon D. Lotter contributed equally to this work and have the right to list their name first in their resumes

This is an open access article under the terms of the Creative Commons Attribution License, which permits use, distribution and reproduction in any medium, provided the original work is properly cited.

© 2023 The Authors. European Eating Disorders Review published by Eating Disorders Association and John Wiley & Sons Ltd.

² WILEY-

Psychosomatics and Psychotherapy, University Hospital RWTH Aachen, Aachen 52074, Germany. Email: lschloesser@ukaachen.de

Funding information

Schweizerische Anorexia Nervosa Stiftung; Bundesministerium für Bildung und Forschung; Max-Planck-Gesellschaft

Highlights

- Sex-specific residual symptoms have been identified in long-term remitted anorexia nervosa (AN) patients, particularly regarding body image disturbance.
- The drive for muscularity (DMS) in male remitted patients suggest a possible risk of transition from male AN to muscle dysmorphic disorder.
- Sex-specific criteria, especially muscle-oriented behaviour, need to be included in the description of diagnoses and symptoms in male patients.

1 | INTRODUCTION AND AIMS

Anorexia nervosa is a severe psychiatric disorder with significant morbidity and mortality that predominantly affects female adolescents and young adults (Herpertz-Dahlmann, 2015). Clinical features include severely restricted food intake leading to or accompanied by low body weight, severe fear of weight gain, and body image disturbance (American Psychiatric Association, 2013). With male-female ratios estimated at approximately 1 to 10-20 and a lifetime prevalence of 0.16%-0.3% (van Eeden et al., 2021), occurrence of AN in boys and men was long considered an "atypical" manifestation of the disorder (Murray, Nagata, et al., 2017). Historically and socially shaped stereotypes, as well as differing clinical presentations, may have further hindered significant improvements in AN diagnosis and treatment in boys and men (Zhang, 2014). Indeed, with the DSM-IV, male patients often did not fulfil the criteria for a "full" AN diagnosis and were diagnosed with an "eating disorder not otherwise specified" (EDNOS), potentially excluding them from both appropriate treatment and contribution to research findings (Raevuori et al., 2014). With the DSM-5, the AN diagnostic criteria, including the amenorrhoea and weight criteria, were modified and the EDNOS category was removed. Given that the numbers of diagnosed male patients have increased since this change (Vo et al., 2016), and considering increasing awareness and improved knowledge of the clinical presentations of AN in males, sex ratios might adapt in future surveys.

To date, the few clinical trials in acute AN that included male patients and explicitly analysed sex differences showed comparable comorbidity profiles, clinical histories, and behavioural symptoms but more pronounced weight and shape concerns in females with AN (Gorrell et al., 2021). While caloric restriction was the most common strategy to achieve weight loss irrespective of sex, male patients seem to engage more strongly in excessive exercising (Silla et al., 2021). The short-term clinical outcome of AN in males was reported to be better than that in females (Coelho et al., 2021). However, the long-term courses were described to be comparable in both sexes, although previous research emphasised the need for a broader range of assessed outcomes in males (e.g., DMS) (Quadflieg et al., 2022). From research findings that were primarily obtained in female patients with AN (Tomba et al., 2019), even in a long-term recovered state and in the absence of any fulfiled DSM-5 criteria, significant residual eating disorder (ED) psychopathology can also be expected for males (Quadflieg et al., 2022). However, the very limited data on potential specific psychopathological and biological characteristics of longterm outcomes of AN in males remitted from AN requires further evaluation.

Consistent with the sex-specific clinical presentations in the form of lower weight phobia and shape concerns but greater excessive exercising in male patients, the body image ideals of patients with acute AN might also differ between sexes. While in females with AN body image disturbance is characterised by an overestimation of one's own body weight and an insatiable desire for thinness (Dalhoff et al., 2019; Provenzano et al., 2019), male patients generally present with a strong DMS and leanness (i.e., low body fat), while low body weight itself is often considered secondary (Gorrell & Murray, 2019). In female patients, as with ED psychopathology, characteristics in perception of body image disturbance often, but not always, improve with weight gain (Dalhoff et al., 2019; Eshkevari et al., 2014). Of note, general population data suggest that muscularity concerns are becoming more prevalent in females as well (Bozsik et al., 2018; Cramblitt & Pritchard, 2013). We therefore emphasise that neither drive for thinness nor DMS is exclusive to either sex or gender, and that more research is needed to delineate the biological and social components of these constructs in birth-assigned males and females as well as in transgender individuals with and without AN. The present study follows this path, although limited to assessing muscle-oriented thoughts and behaviours equivalently in cis-men and -women.

Usually, body image disturbance in AN is assessed using ED-specific interviews and questionnaires, often developed and validated primarily in female populations (e.g., Fairburn et al., 2014). However, muscularity-focused body image ideals, potentially present in males with AN, may be better captured by the *DMS* concept (McCreary & Sasse, 2000), designed as a measure of male body image ideals. Additionally, computer-based body modelling tools have been increasingly applied in AN showing, for example, greater inaccuracy in perception when modelling one's current body and significantly distorted body image ideals if asked to model one's desired body (Mölbert et al., 2018; Ralph-Nearman et al., 2021). While deviant body image ideals are generally well characterised in females with AN, little is known about whether these ideals, and especially their sex-specific variations, occur only in the acute state of the disorder or whether they persist into the long-term course of the disease in both sexes.

Physiologically, eating behaviour is influenced by a variety of hormones that are in turn altered by weight and AN disease status. Indeed, alterations in endocrinological systems can serve as a marker of disease severity and rehabilitation status (Schorr & Miller, 2017). However, data on hormonal status in males with AN, especially in the clinically remitted state, are scarce. During the acute phase of AN, hormonal alterations in free triiodothyronine (fT3), luteinizing hormone (LH), follicle-stimulating hormone (FSH), and leptin, among others, were reported in both men and women (Skolnick et al., 2016; Støving, 2019; Toifl et al., 1988). Compared to healthy controls (HCs) of similar pubertal status, men with acute AN showed lower total testosterone levels (Schorr & Miller, 2017; Silla et al., 2021). In female patients, most endocrine disturbances improve after weight restoration (Baskaran et al., 2017; Schorr & Miller, 2017) and normalise in long-term remission (Wronski et al., 2021). With the exception of a series of three male cases that showed a similar pattern of endocrinological recovery during weight gain to that previously shown in underweight females (Wabitsch et al., 2001), potential hormonal alterations in male patients remitted from AN have not yet been systematically studied.

In this study, we aimed to identify sex-specific residual characteristics of patients with AN in long-term remission using an explorative study design including male and female patients. To quantify residual ED psychopathology and distorted body image, we acquired interview and self-report data and determined aberrant DMS and body image ideals in both males and females with AN using a computer-based 3D modelling tool. Hormonal status was assessed as a marker of biological rehabilitation. Based on the limited literature available to date, we expected significant residual ED psychopathology despite normalised weight in AN patients relative to controls, accompanied by potential sex differences concerning body image ideals.

2 | METHODS

2.1 | Study sample

The study sample consisted of N = 33 patients remitted from AN, n = 24 women (fAN) and n = 9 men (mAN), and 36 HC subjects (fHC, mHC), matched for sex, body mass index (BMI), age, and verbal intelligence (Table 1). "Sex" was defined based on clinical records, and no subject reported a transgender identity. Sexual orientation was assessed via three self-report questions ("To what extent do you consider yourself heterosexual/homosexual/bisexual?"). Patients and male HCs were of central European origin, while 3 female HCs were of Asian origin. All patients (a) had a lifetime history of AN according to DSM-5 criteria (American Psychiatric Association, 2013) and (b) had received inpatient, daycare or outpatient treatment for AN. Healthy control subjects were included if they (a) had a BMI higher than 18 kg/m^2 and (b) had no current or previous psychiatric or neurological disorders as assessed by a screening interview. N = 3 subjects (n = 2 fAN, n = 1mAN) were excluded from the initial cohort of 36 patients because they fulfiled the diagnostic criteria for bulimia nervosa (BN).

To be considered "remitted", patients had to meet the following criteria: (a) no fulfilment of full ED criteria according to the ED Examination (EDE) (Fairburn et al., 2014), (b) a BMI of at least 18 kg/m² or, if underage, at least the 10th percentile, (c) no current treatment for AN, and (d) a stable weight in the time period since the last discharge form inpatient or outpatient treatment. Considering the observed average posttreatment time of 5.5 years (minimum: 1.9 years, maximum: 13.9 years), we adopted the term long-term remitted. According to the DSM-5, partial remission is a sustained period marked by absence of low body weight but persistent core AN symptoms, while full remission is defined as the absence of these criteria (American Psychiatric Association, 2013). Therefore, our criterion (a) includes both definitions for remission.

The study was approved by the local ethics committee (EK 213/15) and was conducted in accordance with the Declaration of Helsinki, and all participants and their legal guardians (if underage) gave written informed consent. Former patients who received inpatient treatment with a diagnosis of AN were recruited from the University Hospital RWTH Aachen, Germany, and surrounding clinics specialising in the treatment of eating disorders. Eligible former patients were identified from the medical records of the department and contacted via mail and/or phone. If interested, a brief telephone screening was conducted (by JO, LDL, or LS), and

TABLE 1 Demographic and clinical sample data.

	Remitted anorexia nervosa (AN)		Healthy controls (HC)		
	Female (N = 24) Mean [SD]/N, Range	Male (N = 9) Mean [SD]/N, Range	Female (N = 26) Mean [SD]/N, Range	Male (N = 10) Mean [SD]/N, Range	Comparison
Age (y)	22.0 [3.2], 17.7-31.4	24.1 [4.4], 18.7–32.5	22.5 [3.5], 16.6–31.3	25.6 [3.7], 20.3–31.0	M > F
Verbal IQ ^a	106.3 [10.4], 85-124	104.8 [12.9], 92–136	110.6 [11.8], 94–130	107.8 [7.2], 95–118	ns.
BMI (kg/m ²)	21.3 [2.7], 18.1–26.6	22.9 [3.5], 18.3–27.4	21.6 [2.1], 18.2–25.5	23.1 [1.9], 19.4–25.5	M > F
BMI-SDS	-0.4 [0.9], -1.7-0.9	-0.4 [1.1], -2.2-0.8	-0.3 [0.7], -1.6-0.9	-0.4 [0.6], -1.4-0.5	ns.
Age at first admission (y) ^b	15.0 [1.5], 12.0–18.0	15.1 [1.9], 12.4–18.1	-	-	ns.
Treatment duration $(y)^b$	1.7 [1.1], 0.1–4.4	2.3 [2.2], 0.1–6.7	-	-	ns.
Post-treatment duration (y) ^b	5.2 [2.7], 1.9–13.9	7.4 [4.5], 2.7–16.3	-	-	ns.
EDE – mean total score	1.0 [0.9], 0.1–3.7	0.5 [0.3], 0.1–1.1	-	-	ns.
EDI-2 – total score ^c	260.1 [70.9], 175–399	213.6 [56.2], 150–321	193.0 [35.8], 151–263	173.3 [21.6], 143–222	$\begin{array}{l} \mathrm{AN} > \mathrm{HC}, \mathrm{W} > \mathrm{M} \\ \mathrm{fAN} > \mathrm{fHC}, \\ \mathrm{mAN} > \mathrm{mHC} \end{array}$
BDI-2 ^d	11.3 [12.0], 0–37	7.8 [8.5], 0–26	5.2 [4.7], 0–14	3.3 [3.7], 0–10	AN > HC, fAN > fHC
BAI ^e	6.7 [7.1], 0–29	4.8 [4.3], 2–13	4.4 [5.8], 0-24	1.3 [1.0], 0–3	ns.
SF-36 – total score ^f	84.6 [14.2], 42.5–114	84.7 [8.4], 68.6–94.6	83.9 [7.9], 62.5–95.4	91.4 [4.0], 84.9–96.4	ns.
SF-36 – mental health ^g	79.6 [17.4], 35–100	71.9 [22.0], 35–95	80.2 [13.4], 50-100	85.9 [10.8], 72–100	ns.
Heterosexuality	6.7 [1.0], 2–7	7 [0], 7	6.1 [1.8], 1–7	6.9 [0.3], 6-7	ns.
Homosexuality	1 [0], 1	1 [0], 1	1.0 [0.2], 1–2	1 [0], 1	Ns.
Bisexuality	1.2 [0.4], 1–2	1 [0], 1	2.1 [1.6], 1-6	1.1 [0.3], 1–2	
Hormonal contraception	10	-	19	-	fHC > fAN
MINI – ≥ 1 comorbidity	8	1	-	-	ns.

Note: Variables were compared between groups as appropriate, using two-way analyses of variances with bootstrapping followed by evaluation of simple main effects, Mann-Whitney U tests, or Chi-squared tests. Missing values: a: n = 1 fHC, 1 mHC. b: n = 1 fAN, 1 mAN. c: n = 1 fHC. d: n = 1 fAN, 1 mAN, 3 fHC, 1 mHC. e: n = 3 fAN, 1 mAN, 5 fHC, 4 mHC. f: n = 2 fAN, 1 mAN, 1 fHC, 2 mHC. g: n = 1 fAN, 1 mAN, 1 fHC, 2 mHC. SD = standard deviation, y = years, IQ = intelligence quotient, BMI (-SDS) = body mass index (-standard deviation score), EDE = Eating Disorder Examination, EDI-2 = Eating Disorder Inventory, BDI-2 = Beck Depression Inventory 2, BAI = Beck Anxiety Inventory, MINI = Mini International Neuropsychiatric Interview, M/m = male, F/f = female, ns. = not significant.

participants who reported no current ED symptoms and/ or ongoing treatment were invited to the department for validation of their remission status and further study assessments. Healthy control subjects were recruited via postings from the surroundings of the University Hospital RWTH Aachen without being informed about the goals of the study. All participants received financial compensation in accordance with the local standards for clinical studies.

The data and results reported here were published in part in LS's medical thesis (Schlösser et al. (2022); neuroimaging data collected from an overlapping sample (n = 21 female patients and N = 22 female HCs) were published in another report (Lotter et al., 2021).

2.2 | Clinical characteristics and psychopathology

In patients, data on the current state and long-term clinical course of EDs were collected in structured interviews and from medical records if the latter were available (n = 31 patients). *Treatment duration* was defined as the time between first admission and last

discharge from inpatient and/or outpatient treatment with a diagnosis of AN; posttreatment duration was defined as the time between last discharge and the study date. Body weight was measured with a standardized scale in the morning prior to food intake and adjusted for height, age, and sex by calculation of the BMIstandard deviation score (BMI-SDS) (Hemmelmann et al., 2010; Neuhauser et al., 2013). Eating disorder diagnostic criteria and severity were assessed using the Eating Disorder Examination (EDE) and the Eating Disorder Inventory 2 (EDI) (Fairburn et al., 2014; Garner, 1991). Eating Disorder Examination criteria for diagnosis of AN and BN according to DSM-5 were selected to assess weight phobia (AN-B: Fear of gaining weight), body image disturbance (AN-C: Disturbance of body weight/shape experience), binge eating (BN-A: *Recurrent* episodes of binge eating), and "purging" behaviour (BN-B: Recurrent inappropriate compensatory behaviours to prevent weight gain). Comorbid psychiatric disorders were screened using the Mini International Neuropsychiatric Interview (MINI) (Ackenheil et al., 1999). The Beck Anxiety Inventory (BAI) (Margraf & Ehlers, 2007) and the revised Beck Depression Inventory (BDI) (Hautzinger et al., 2006) were used to assess anxiety and depressive symptoms. Health-related quality of life was assessed with the 36-Item Short Form Health Survey (SF-36) (Lins & Carvalho, 2016). The Multiple-choice Vocabulary Test (Lehrl, 2005) was utilised to approximate the verbal intelligence quotient. Interviews were conducted by a specifically trained and supervised clinical researcher (JO).

2.3 | Body image and drive for muscularity

Characteristics of body image disorder were assessed using 3D modelling software ("Body Morphing Tool" (BMT), DAZ Studio 4 Pro, Daz3D.com) (Crossley et al., 2012). Participants were seated in front of a 15-inch notebook and asked to model their own body (a) as they perceived it at the time of testing and (b) as they imagined their "desired" body. The avatars were sex-specific and could be modified by slide controls in five categories selected from the software's presets: "Body Size", "Weight" (ranging from -100 to +100%), "Bodybuilder", "Fitness", and "Stocky" in males or "Voluptuous" in females (ranging from 0% to 100%). The BMT allowed free rotation and zooming into the body (Figure S1). Values corresponding to the slide control positions of the categories "Body builder", "Fitness", and "Weight" were used for group comparisons (hereafter referred to as muscu*larity, definition, and body fat).* From the five groupwise mean BMT indices, we created visualizations of the "average" perceived and desired bodies for each group. To assess muscularity-related attitudes and behaviour, we applied the Drive for Muscularity Scale (DMS), consisting of 15 questions (Waldorf et al., 2014).

2.4 | Hormonal status

Blood was drawn in the morning after an overnight fast, centrifuged, aliquoted, and stored at -80 °C. Concentrations of leptin, fT3, cortisol, LH, FSH, progesterone, oestradiol, and testosterone were determined by enzyme-linked immunosorbent assays (laboratory of the University Hospital Magdeburg, Germany). Subjects using hormonal contraception were excluded from the analyses of FSH, LH, progesterone, oestrogen, and testosterone because of its influence on sex hormone levels.

2.5 | Statistical analyses and visualizations

Statistical tests were carried out using SPSS (IBM Corp., 2020, version 26). Due to multiple nonnormally distributed variables as evaluated using Shapiro-Wilk tests and the small sample sizes of the male groups, nonparametric statistics and bootstrapping procedures were used whenever possible. Because of the small group sizes, EDE diagnostic criteria were compared between groups using two-sided Fisher's exact tests. Demographic, clinical, and self-report data (EDI subscales, EDE, BAI, BDI, SF-36, DMS), hormonal plasma concentrations, and parameters derived from the BMT were compared using two-way analyses of covariance (ANCOVAs) with diagnosis and sex as main factors. Age and BMI-SDS were included as covariates in all models. Main and diagnosis \times sex interaction effects were evaluated and followed up on by evaluation of the simple main effects of diagnosis within sex categories (Šidák correction). Bootstrapping procedures as implemented in SPSS (N = 1000 samples; random number generator: Mersenne Twister, seed: 2,000,000) were used in ANCOVA models to account for violations of normality assumptions (Chernick, 2008).

In sensitivity analyses, we repeated all ANCOVAs while (a) applying the 10th-percentile BMI threshold to all subjects irrespective of age, (b) including three additional subjects with current BN, (c) applying binary logistic regression to control significant group comparisons of EDE criteria for the influence of BMI-SDS and age, and (d) using Mann–Whitney U (MWU) tests to assess significant ANCOVA results for the potential impact of

outliers. Correlational analyses were conducted separately within the respective subgroups using Spearman correlations. Due to the exploratory nature of this study and the assessment of multiple intercorrelated variables, the significance level was set at two-tailed $\alpha = 0.05$ and Bonferroni-corrected per set of analyses (in the case of ANCOVAs, applied to assess the significance of the complete model): EDE criteria (4 tests), EDI subscales (11 tests), hormone levels (8 tests), and DMS/BMT (7 tests). Given the early state of the research on sex differences in AN, the literature was insufficient to conduct a priori power analyses.

Visualizations were created using ggplot2, version 3.3.5 (Wickham, 2016), in R, version 3.6.3 (R Core Team, 2021).

3 | RESULTS

3.1 | Sample characteristics and general results

Age differed between males and females but not between former patients and HCs. No group differences were found for verbal intelligence, BMI, or BMI-SDS. Sexual orientation was largely heterosexual and did not differ between groups (n = 2 fHC: bisexual orientation, n = 1fHC: no sexual preference). Female and male patients were comparable regarding age at admission, treatment and posttreatment durations, EDE total scores, and current number of comorbidities. Patients presented with higher overall ED, depression, and anxiety scores than HCs, but the groups did not differ regarding quality of life (Table 1, Table S1).

3.2 | Residual eating disorder psychopathology

applying Bonferroni correction (corrected When $\alpha = 0.0125$), EDE criteria frequency did not differ between male and female patients remitted from AN. However, numerically, more male than female patients fulfiled the EDE criterion BN-B: Recurrent inappropriate compensatory behaviours to prevent weight gain, although the result was not significant ($\chi^2(1) = 7.64$, p = 0.013, uncorrected), while the opposite was the case for AN-C: Disturbance of body weight/shape experience ($\chi^2(1) = 4.91$, p = 0.044, uncorrected), being observed more frequently in females. Further analyses revealed that compensatory behaviours consisted of pathologically driven exercising in all mAN subjects. The other criteria did not differ between the sex groups (AN-B: $\chi^2(1) = 0.39$, p = 1.000;

BN-A: $\chi^2(1) = 0.76$, p = 0.642; Figure 1a). Bootstrapped binary logistic regression analyses of criteria BN-B and AN-C indicated robustness against influences of age and BMI-SDS (BN-B: $\chi^2(3) = 7.89$, p = 0.048, Sex: B = -2.39, SE = 7.27, p = 0.010; AN-C: $\chi^2(3) = 5.11$, p = 0.164, Sex: B = -1.94, SE = 6.92, p = 0.013).

Bootstrapped two-way ANCOVA models of the EDI subscales were significant for the scales drive for thinness, body dissatisfaction, ineffectiveness, interoceptive awareness, and impulse regulation. In all cases, we found significant main effects of diagnosis, with AN patients showing higher scores than HCs with large effect sizes. A significant main effect of sex revealed a higher drive for thinness and body dissatisfaction in females than in males. The analyses of simple main effects further indicated that the effects of diagnosis were strongly driven by the group differences between fAN patients and fHCs; however, no interaction effects emerged. When applying Bonferroni correction, no model was significant, but some main effects emerged for the remaining subscales (Table S3, Figure 1b). Recalculating the significant models while (a) excluding participants with a BMI below the 10th percentile, (b) including subjects with current BN, and (c) using MWU tests confirmed the stability of the group effects (Tables S4-S6).

3.3 | Body image and drive for muscularity

ANCOVAs assessing the BMT-derived variables desired muscularity and desired and perceived muscle definition were significant on the model level and showed significant sex, diagnosis \times sex interaction, and, except for desired muscle definition, diagnosis effects (Bonferroni corrected, large effect sizes). For perceived muscularity, we observed effects of the diagnosis main factor and the interaction (uncorrected). Body fat indices did not differ between groups. Regarding self-reported DMS, significant effects of diagnosis and sex emerged. In all cases, AN patients exhibited increased scores relative to HCs, and males showed higher scores than females. Simple main effects indicated a general pattern of higher DMS in mAN patients than in mHCs that was not present in female subgroups (Table S3, Figure 2a). Group difference patterns largely remained identical when recalculating analyses while applying the stricter BMI criterion, including BN subjects, or using MWU tests (Tables S4-S6). Figure 2b shows groupwise "average" avatars for the perceived and desired conditions. As measures of potential clinical value, we additionally evaluated group differences and clinical correlations of BMT "difference" metrics, calculated as



female HC i female AN i male AN i male HC

FIGURE 1 Residual eating disorder (ED) psychopathology and hormonal status. (a) Percentages of female and male patients fulfiling the preselected diagnostic criteria for anorexia nervosa (AN) (AN-B: *Fear of gaining weight*, AN-C: *Disturbance of body weight/shape experience*) and bulimia nervosa (BN) (BN-A: *Recurrent episodes of binge eating*, BN-B: *Recurrent inappropriate compensatory behaviours to prevent weight gain*). Numbers in bars: "percent (n)". (b) and (c) Group-wise results of EDI subscales and hormone assessments, respectively. Hormone levels are given per millilitre and log10-scaled. Individual values are shown as scatter points and squares, violins depict distributions of values within each group (not scaled to group size but to same width per group), red circles and error bars show group-wise mean values and corresponding bias-corrected and accelerated (BCa) bootstrap confidence intervals. Brackets show significant group comparison results, **p* < 0.05, ***p* < 0.01, ****p* < 0.001, D = effect of diagnosis, S = effect of sex, DxS = interaction effect. AN = Anorexia nervosa, BN = Bulimia nervosa, HC = Healthy control, EDI = Eating Disorder Inventory 2, fT3 = free triiodothyronine, LH = luteinizing hormone, FSH = follicle-stimulating hormone. See Tables S2 and S3 for detailed results.

the subject-wise difference between each perceived and desired condition. We found that males independent of diagnosis had a higher discrepancy between their *desired* and *perceived muscularity* and *fitness* (Table S7, Figure S2).

3.4 | Hormonal status

ANCOVA models were significant only for leptin, cortisol, and testosterone (Bonferroni corrected; Table S3, Figure 1c). In all cases, plasma levels were significantly affected by sex, with females showing higher levels of leptin and cortisol and males demonstrating higher testosterone levels. For testosterone, we additionally observed a significant interaction between diagnosis and sex with a medium effect size. This effect emerged from reduced testosterone levels in the male AN group but increased testosterone levels in the female

AN group compared to sex-specific HC groups. Note, however, that the simple main effects were not significant and that the effect size for the interaction effect was medium. Sensitivity analyses showed no relevant influence of a more conservative BMI cut-off or the additional inclusion of BN subjects on all group effects and a general stability of results against influence of outliers (Tables S4–S6).

3.5 | Correlation analyses

Leptin correlated strongly with BMI-SDS in female subgroups but not in males; testosterone and BMI did not show significant associations in any subgroup (Figure S3). Evaluating associations between BMT indices, DMS, *drive for thinness*, and body composition, we found (a) patterns of positive associations between *perceived body fat* and BMI-SDS as well as between leptin





FIGURE 2 Body image characteristics. (a) Group-wise results of drive for muscularity (DMS) Scale and Body Morphing Tool (BMT) parameters. See Figure 1 for descriptions. (b) Group-wise averaged perceived and desired avatars, derived by manually configuring avatars from mean values of five BMT scales and rendering within *DAZ Studio 4 Pro*.

and *drive for thinness*, especially in female subgroups, (b) negative associations between *desired body fat* and DMS scores across groups, and (c) a strong correlation between perceived muscularity and BMI-SDS only in males with AN (Table S8, Figure S4). Exploratory correlational analyses with BMT difference metrics revealed a consistent pattern of strong negative discrepancies between *desired* and *perceived body fat* (i.e., to desire less body fat than

one perceives to have) in females with high DMS, thinness, and total EDI scores. The opposite pattern was observed between *muscularity* and the self-report scales (i.e., higher ED scores in subjects who desire more muscularity than they perceive to have). Interestingly, the latter relationships were observed in both male groups and in females remitted from AN, but not in healthy females (Figure S5).

4 | DISCUSSION

In this exploratory study, we investigated ED psychopathology, body image ideals, and hormonal status in female and male patients with long-term remission from AN. We found significantly elevated ED psychopathology focused on body dissatisfaction in females as opposed to excessive exercising in males. In line with this finding, male patients remitted from AN pursued significantly muscularity-centred body image ideals compared to females and male controls.

Despite the persistent ED symptomatology, we observed almost full endocrinological rehabilitation in remitted AN which was most likely achieved by regular and sufficient energy intake (Støving, 2019). The observed sex differences in leptin, testosterone, and cortisol levels can be considered physiological (Asarian & Geary, 2013; Christen et al., 2018; Goel et al., 2014). Elevated ED psychopathology in (female) patients in various states of recovery from AN can be considered a well-replicated finding (Bernardoni et al., 2020; Boehm et al., 2016; Cascino et al., 2020), and thus, our results were to be expected. Eating disorder psychopathology differing by sex, with females showing stronger concerns about body weight and shape and males exhibiting especially pronounced driven exercising, was previously reported independently in females and males with acute (Gravina et al., 2018; Mairs & Nicholls, 2016) and shortterm remitted AN (Heimann et al., 2018; Silla et al., 2021). We extend these results by showing similar patterns in long-term remission of AN in the form of significantly elevated body dissatisfaction (EDI-2) and a higher frequency of body image disturbance in female remitted AN patients but a significantly higher frequency of compensatory behaviours in male remitted AN patients (EDE).

In line with the above findings, using the BMT, we demonstrated body image ideals centred on muscularity and fitness, accompanied by increased muscularityoriented attitudes and behaviours (DMS), in male patients remitted from AN in a more naturalistic manner, as was previously reported in acute AN (Murray et al., 2016). In contrast, female patients did not differ from controls on muscularity and body fat ratings, irrespective of the *perceived* or *desired* conditions. On the one hand, this could be interpreted as evidence for a normalisation of the distorted body image in females remitted from AN. On the other hand, especially considering the persistently elevated ED psychopathology and DMS observed in the same patients and the interaction effect of diagnosis and sex on BMT scores, a potential repulsive effect of higher muscularity and subsequently higher body weight specifically in female

patients might be discussed. Given the preliminary nature of this study, these findings require replication in a larger cohort based on tools validated in both female and male populations. Relatedly, we found that the BMT seemed to capture individual discrepancies between one's ideal and one's perceived body, indicating its potential value as a diagnostic tool to evaluated in future studies.

Considering that all male patients in this study initially had low body weight and were diagnosed with AN, the observed muscularity-focused body image ideals suggest a possible risk of transition to muscle dysmorphic disorder in the long-term clinical course of males with AN. In clinical practice, it should be considered that a comorbid muscle dysmorphic disorder may have already been present at the first clinical presentation. Muscle dysmorphic disorder, also referred to as "reverse anorexia", is characterised by the desire for a larger and more muscular body accompanied by the belief that one is small and thin, even though individuals are often tall and muscular (Murray et al., 2010). Indeed, recently, a reconsideration of muscle dysmorphic disorder from the perspective of ED pathology (Gorrell & Murray, 2019) and the corresponding adaptation of treatment strategies (Cunningham et al., 2017) have been proposed. Transitions from AN to muscle dysmorphic disorder in male patients have been previously described (Murray, Griffiths, et al., 2017). However, whether these muscle dysmorphic body image distortions actually have negative impacts on patients' quality of life remains unclear and could not be verified by the present investigation. However, the association between AN and muscle dysmorphic disorder in males calls for further investigation.

4.1 | Limitations of the present study and considerations for future research

This exploratory study is subject to several limitations. However, given the clinical relevance of the topic at hand, we consider our findings an important first step and a blueprint for future research projects to learn from and build upon. First, this exploratory study is limited by the small sample size, especially of the male subgroup, requiring replication in well-powered studies. Second, it is questionable whether measurements and tests developed primarily for female patients with AN adequately assess AN diagnosis and symptoms in males. The EDE appears to have clinical utility in men (Schaefer et al., 2019), although they tend to score lower than females (Darcy et al., 2012). On the other hand, the EDI-2 did not detect sex differences in the present study, which is in line with its previously reported limited reliability in men (Spillane et al., 2004). Third, the usually applied

weight criterion of a threshold BMI of 18.5 kg/m² corresponds to the 10th BMI percentile only in females, but not in males, which generally complicates comparison of sexes in AN. Here, we further included participants based on a BMI cut-off of at least 18 kg/m², whereas the DSM-5 defines "significantly low body-weight" with a BMI threshold of 18.5 kg/m². However, our sensitivity analyses using a more conservative BMI cut-off largely confirmed our findings. In line with the DSM approach, defining "recovery" based on a weight cutoff alone has proven insufficient (Khalsa et al., 2017), suggesting that a comprehensive definition of recovery must include a psychological dimension (Bardone-Cone et al., 2018). Given the long average posttreatment time of 5.5 years but still persisting psychopathology, we decided to classify our former patient cohort as "in remission" rather than as "recovered". Whether it is now useful to distinguish between "remission" and "recovery" based solely on post-treatment duration, as has often been done in similar studies, should be investigated in future research. Fourth, our approach to measuring body image characteristics is limited, as it only assessed self-reported muscularity-oriented cognition and behaviour as well as body image ideals but lacked assessment of actual fat or muscle mass to validate the subjective measures. In addition, the BMT template avatars were white and not individualised, which could limit the validity of the approach, as participants may not identify with the avatars.

5 | CONCLUSION

In line with previous studies (Bardone-Cone et al., 2019), we found evidence for sex-specific characteristics of body image disturbance and DMS in long-term remitted AN indicating a potential association between male AN and muscle dysmorphic disorder. Thus, we recommend addressing sex-specific ED symptomatology, in particular muscularity-oriented body image ideals and behaviours when assessing diagnosis and symptoms in males with AN. Incorporation of sex-specific symptoms, as well as body weight (BMI) thresholds, in AN criteria for diagnosis, remission, and recovery will have to be discussed in the future. This will require further development of appropriate assessment instruments, especially for male patients with AN (Arkenau et al., 2020).

AUTHOR CONTRIBUTIONS

LS and LDL performed data analyses and wrote the manuscript. KK, JS and BHD designed the overall study. LS, LDL and JO conducted the clinical study and gathered the data under supervision of JS, KK and BHD. KB and RB monitored the hormone assessments. All authors reviewed and commented on the manuscript.

ACKNOWLEDGEMENTS

We thank the staff of the DRK Fachklinik Bad-Neuenahr, the University Medical Center Utrecht, and the University Hospital Vienna for their support in providing patient contacts and clinical data. LDL was supported by the Federal Ministry of Education and Research (BMBF) and the Max Planck Society (MPG), Germany.

The study was supported by the Swiss AN foundation (Grant number: 53-15).

Open Access funding enabled and organized by Projekt DEAL.

CONFLICT OF INTEREST STATEMENT

No author reports conflicts of interest with respect to the work presented in this study.

DATA AVAILABILITY STATEMENT

The data presented in the study are available upon reasonable request from the author.

ORCID

Louisa Schloesser D https://orcid.org/0000-0002-8134-7377

Leon D. Lotter b https://orcid.org/0000-0002-2337-6073 Jochen Seitz b https://orcid.org/0000-0002-0110-7980 Kerstin Konrad b https://orcid.org/0000-0001-9039-2615 Beate Herpertz-Dahlmann b https://orcid.org/0000-0001-8450-3323

REFERENCES

- Ackenheil, M., Stotz, G., & Dietz-Bauer, R. (1999). Mini international neuropsychiatric interview. German version 5.0.0, DSM-IV. Psychiatrische Universitätsklinik München.
- American Psychiatric Association. (2013). Diagnostic and statistical manual of mental disorders (DSM-5^{*}). Retrieved from https:// books.google.de/books?id=-JivBAAAQBAJ
- Arkenau, R., Vocks, S., Taube, C. O., Waldorf, M., & Hartmann, A. S. (2020). The Body Image Matrix of Thinness and Muscularity-Male Bodies: Development and validation of a new figure rating scale for body image in men. *Journal of Clinical Psychology*, 76(7), 1283–1292. https://doi.org/10.1002/ jclp.22933
- Asarian, L., & Geary, N. (2013). Sex differences in the physiology of eating. Am J physiol regul integr. *Comparative Physiology*, 305(11), R1215–R1267. https://doi.org/10.1152/ajpregu.00446. 2012
- Bardone-Cone, A. M., Hunt, R. A., & Watson, H. J. (2018). An overview of conceptualizations of eating disorder recovery, recent findings, and future directions. *Current Psychiatry Reports*, 20(9), 79. https://doi.org/10.1007/s11920-018-0932-9
- Bardone-Cone, A. M., Johnson, S., Raney, T. J., Zucker, N., Watson, H. J., & Bulik, C. M. (2019). Eating disorder recovery in men: A

pilot study. International Journal of Eating Disorders, 52(12), 1370–1379. Epub 2019 Aug 16. PMID: 31418898. https://doi. org/10.1002/eat.23153

- Baskaran, C., Misra, M., & Klibanski, A. (2017). Effects of anorexia nervosa on the endocrine system. *Pediatric Endocrinology Reviews*, 14(3), 302–311. https://doi.org/10.17458/per.vol14.2017. BMK.effectsanorexianervosa
- Bernardoni, F., Bernhardt, N., Pooseh, S., King, J. A., Geisler, D., Ritschel, F., Boehm, I., Seidel, M., Roessner, V., Smolka, M. N., & Ehrlich, S. (2020). Metabolic state and value-based decision-making in acute and recovered female patients with anorexia nervosa. *Journal of Psychiatry & Neuroscience*, 45(4), 253–261. https://doi.org/10.1503/jpn.190031
- Boehm, I., Finke, B., Tam, F. I., Fittig, E., Scholz, M., Gantchev, K., Roessner, V., & Ehrlich, S. (2016). Effects of perceptual body image distortion and early weight gain on long-term outcome of adolescent anorexia nervosa. *European Child & Adolescent Psychiatry*, 25(12), 1319–1326. https://doi.org/10.1007/s00787-016-0854-1
- Bozsik, F., Whisenhunt, B. L., Hudson, D. L., Bennett, B., & Lundgren, J. D. (2018). Thin is in? Think again: The rising importance of muscularity in the thin ideal female body. *Sex Roles*, 79(9), 609–615. https://doi.org/10.1007/s11199-017-0886-0
- Cascino, G., Canna, A., Monteleone, A. M., Russo, A. G., Prinster, A., Aiello, M., Esposito, F., Salle, F. D., & Monteleone, P. (2020). Cortical thickness, local gyrification index and fractal dimensionality in people with acute and recovered Anorexia Nervosa and in people with Bulimia Nervosa. *Psychiatry Research Neuroimaging*, 299, 111069. https://doi.org/10.1016/j. pscychresns.2020.111069
- Chernick, M. R. (2008). Bootstrap methods: A guide for practitioners and researchers. 2. Auflage. Wiley.
- Christen, T., Trompet, S., Noordam, R., van Klinken, J. B., van Dijk, K. W., Lamb, H. J., Cobbaert, C. M., den Heijer, M., Jazet, I. M., Jukema, J. W., Rosendaal, F. R., & de Mutsert, R. (2018). Sex differences in body fat distribution are related to sex differences in serum leptin and adiponectin. *Peptides*, 107, 25–31. https://doi.org/10.1016/j.peptides.2018. 07.008
- Coelho, J. S., Suen, J., Marshall, S., Burns, A., Geller, J., & Lam, P. Y. (2021). Sep 15). Gender differences in symptom presentation and treatment outcome in children and youths with eating disorders. *J Eat Disord*, 9(1), 113. https://doi.org/10. 1186/s40337-021-00468-8
- Cramblitt, B., & Pritchard, M. (2013). Media's influence on the drive for muscularity in undergraduates. *Eating Behaviors*, 14(4), 441–446. https://doi.org/10.1016/j.eatbeh.2013.08.003
- Crossley, K. L., Cornelissen, P. L., & Tovée, M. J. (2012). What is an attractive body? Using an interactive 3D program to create the ideal body for you and your partner. *PLoS One*, *7*(11), e50601. https://doi.org/10.1371/journal.pone.0050601
- Cunningham, M. L., Griffiths, S., Mitchison, D., Mond, J. M., Castle, D., & Murray, S. B. (2017). Muscle dysmorphia: An overview of clinical features and treatment options. *Journal of Cognitive Psychotherapy*, 31(4), 255–271. https://doi.org/10. 1891/0889-8391.31.4.255
- Dalhoff, A. W., Romero Frausto, H., Romer, G., & Wessing, I. (2019). Perceptive body image distortion in adolescent

WILEY 11

anorexia nervosa: Changes after treatment. Frontiers in Psychiatry, 10, 748. https://doi.org/10.3389/fpsyt.2019.00748

- Darcy, A. M., Doyle, A. C., Lock, J., Peebles, R., Doyle, P., & Le Grange, D. (2012). The eating disorders examination in adolescent males with anorexia nervosa: How does it compare to adolescent females? *International Journal of Eating Disorders*, 45(1), 110–114. https://doi.org/10.1002/eat.20896
- Eshkevari, E., Rieger, E., Longo, M. R., Haggard, P., & Treasure, J. (2014). Persistent body image disturbance following recovery from eating disorders. *International Journal of Eating Disorders*, 47(4), 400–409. https://doi.org/10.1002/eat.22219
- Fairburn, C. G., Cooper, Z., & O'Connor, M. E. (2014). Eating disorder examination (edition 17.0D). Retrieved from http:// www.credo-oxford.com/7.2.htm
- Garner, D. M. (1991). Eating disorder inventory-2: Professional manual. Psychological Assessment Resources, 0–70.
- Goel, N., Workman, J. L., Lee, T. T., Innala, L., & Viau, V. (2014). Sex differences in the HPA axis. *Comprehensive Physiology*, 4(3), 1121–1155. https://doi.org/10.1002/cphy.c130054
- Gorrell, S., Hughes, E. K., Sawyer, S. M., Roberts, S. R., Nagata, J. M., Yeo, M., Lock, J., & Le Grange, D. (2021). Gender-based clinical differences in evidence-based treatment for adolescent anorexia nervosa: Analysis of aggregated randomized controlled trials. *Eating and Weight Disorders*, 27(3), 1123–1130. https://doi.org/10.1007/s40519-021-01257-7
- Gorrell, S., & Murray, S. B. (2019). Eating disorders in males. Child Adolesc Psychiatr Clin N Am, 28(4), 641–651. https://doi.org/ 10.1016/j.chc.2019.05.012
- Gravina, G., Milano, W., Nebbiai, G., Piccione, C., & Capasso, A. (2018). Medical complications in anorexia and bulimia nervosa. Endocrine, Metabolic & amp; Immune Disorders: Drug Targets, 18(5), 477–488. https://doi.org/10.2174/1871530318 666180531094508
- Hautzinger, M., Keller, F., & Kuhner, C. (2006). Das Beck Depressions-inventar II. Deutsche Bearbeitung und Handbuch zum BDI II. Harcourt Test Services, Frankfurt a. M.
- Heimann, P. M., Konrad, K., & Vloet, T. D. (2018). Anorexia nervosa in males. Zeitschrift f
 ür Kinder- und Jugendpsychiatrie und Psychotherapie, 46(6), 478–487. (Anorexia nervosa bei Jungen). https://doi.org/10.1024/1422-4917/a000579
- Hemmelmann, C., Brose, S., Vens, M., Hebebrand, J., & Ziegler, A. (2010). [Percentiles of body mass index of 18-80-year-old German adults based on data from the Second National Nutrition Survey]. *Deutsche medizinische Wochenschrift, 135*(17), 848–852. (Perzentilen des Body-Mass-Index auch für 18- bis 80-Jährige? Daten der Nationalen Verzehrsstudie II). https://doi.org/10.1055/s-0030-1253666
- Herpertz-Dahlmann, B. (2015). Adolescent eating disorders: Update on definitions, symptomatology, epidemiology, and comorbidity. *Child Adolesc Psychiatr Clin N Am*, 24(1), 177–196. https://doi.org/10.1016/j.chc.2014.08.003
- Khalsa, S. S., Portnoff, L. C., McCurdy-McKinnon, D., & Feusner, J. D. (2017). What happens after treatment? A systematic review of relapse, remission, and recovery in anorexia nervosa. J Eat Disord, 5(1), 20. https://doi.org/10.1186/s40337-017-0145-3
- Lehrl, S. (2005). Mehrfachwahl-wortschatz-intelligenztest: MWT-B (5., unveränd. Aufl. Ed.). Spitta.
- Lins, L., & Carvalho, F. M. (2016). SF-36 total score as a single measure of health-related quality of life: Scoping review. *SAGE*

10990968, 0, Downloaded from https://onlinelibrary.wiley.com/doi/10.1002/erv.2997 by Cochrane Germany, Wiley Online Library on [16/06/2023]. See the Terms and Conditions (https://onlinelibrary.wiley.com/terms-and-conditions) on Wiley Online Library for rules of use; OA articles are governed by the applicable Creative Commons License

Open Medicine, 4, 2050312116671725. https://doi.org/10.1177/ 2050312116671725

- Lotter, L. D., von Polier, G., Offermann, J., Buettgen, K., Stanetzky,
 L., Eickhoff, S. B., Konrad, K., Seitz, J., & Dukart, J. (2021).
 Recovery-associated resting-state activity and connectivity alterations in anorexia nervosa. *Biol Psychiatry Cogn Neurosci Neuroimaging*, 6(10), 1023–1033. Epub 2021 Mar 23. PMID: 33766777. https://doi.org/10.1016/j.bpsc.2021.03.006
- Mairs, R., & Nicholls, D. (2016). Assessment and treatment of eating disorders in children and adolescents. Archives of Disease in Childhood, 101(12), 1168–1175. https://doi.org/10.1136/ archdischild-2015-309481
- Margraf, J., & Ehlers, A. (2007). *Beck angstinventar deutsche version* (*Bai*). Hans Huber.
- McCreary, D. R., & Sasse, D. K. (2000). An exploration of the drive for muscularity in adolescent boys and girls. *Journal of American College Health*, 48(6), 297–304. https://doi.org/10. 1080/07448480009596271
- Mölbert, S. C., Thaler, A., Mohler, B. J., Streuber, S., Romero, J., Black, M. J., Zipfel, S., Karnath, H. O., & Giel, K. E. (2018). Assessing body image in anorexia nervosa using biometric selfavatars in virtual reality: Attitudinal components rather than visual body size estimation are distorted. *Psychological Medicine*, 48(4), 642–653. https://doi.org/10.1017/s00332917170 02008
- Murray, S. B., Griffiths, S., & Mond, J. M. (2016). Evolving eating disorder psychopathology: Conceptualising muscularityoriented disordered eating. *British Journal of Psychiatry*, 208(5), 414–415. https://doi.org/10.1192/bjp.bp.115.168427
- Murray, S. B., Griffiths, S., Mitchison, D., & Mond, J. M. (2017). The transition from thinness-oriented to muscularity-oriented disordered eating in adolescent males: A clinical observation. *Journal of Adolescent Health*, 60(3), 353–355. https://doi.org/ 10.1016/j.jadohealth.2016.10.014
- Murray, S. B., Nagata, J. M., Griffiths, S., Calzo, J. P., Brown, T. A., Mitchison, D., Blashill, A. J., & Mond, J. M. (2017). The enigma of male eating disorders: A critical review and synthesis. *Clinical Psychology Review*, 57, 1–11. https://doi.org/10. 1016/j.cpr.2017.08.001
- Murray, S. B., Rieger, E., Touyz, S. W., & De la Garza García Lic, Y. (2010). Muscle dysmorphia and the DSM-V conundrum: Where does it belong? A review paper. *International Journal of Eating Disorders*, 43(6), 483–491. https://doi.org/10.1002/eat. 20828
- Neuhauser, H., Schienkiewitz, A., Rosario, A. S., Dortschy, R., & Kurth, B.-M. (2013). Referenzperzentile für anthropometrische Maßzahlen und Blutdruck aus der Studie zur Gesundheit von Kindern und Jugendlichen in Deutschland (KiGGS). https:// doi.org/10.25646/3179
- Provenzano, L., Porciello, G., Ciccarone, S., Lenggenhager, B., Tieri, G., Marucci, M., Dazzi, F., Loriedo, C., & Bufalari, I. (2019). Characterizing body image distortion and bodily self-plasticity in anorexia nervosa via visuo-tactile stimulation in virtual reality. *Journal of Clinical Medicine*, 9(1), 98. https://doi.org/10. 3390/jcm9010098
- Quadflieg, N., Naab, S., Voderholzer, U., & Fichter, M. M. (2022). Long-term outcome in males with anorexia nervosa: A prospective, sex-matched study. *International Journal of Eating Disorders*, 55(3), 393–398. https://doi.org/10.1002/eat.23672

- Raevuori, A., Keski-Rahkonen, A., & Hoek, H. W. (2014). A review of eating disorders in males. *Current Opinion in Psychiatry*, *27*(6), 426–430. https://doi.org/10.1097/yco.00000000 0000113
- Ralph-Nearman, C., Arevian, A. C., Moseman, S., Sinik, M., Chappelle, S., Feusner, J. D., & Khalsa, S. S. (2021). Visual mapping of body image disturbance in anorexia nervosa reveals objective markers of illness severity. *Scientific Reports*, *11*(1), 12262. https://doi.org/10.1038/s41598-021-90739-w
- R Core Team. (2021). R: A language and environment for statistical computing. Retrieved from https://www.R-project.org/}
- Schaefer, L. M., Anderson, L. M., Simone, M., O'Connor, S. M., Zickgraf, H., Anderson, D. A., Rodgers, R. F., & Thompson, J. K. (2019). Gender-based differential item functioning in measures of eating pathology. *International Journal of Eating Disorders*, 52(9), 1047–1051. https://doi.org/10.1002/ eat.23126
- Schlösser, L., Konrad, K., & Habel, U. (2022). Klinischer und endokrinologischer Status bei langzeitgewichtsrehabilitierten Patientinnen und Patienten mit Anorexia nervosa. [Dissertation / PhD Thesis, RWTH Aachen University]. Retrieved from https://publications.rwth-aachen.de/record/849848
- Schorr, M., & Miller, K. K. (2017). The endocrine manifestations of anorexia nervosa: Mechanisms and management. *Nature Reviews Endocrinology*, 13(3), 174–186. https://doi.org/10.1038/ nrendo.2016.175
- Silla, K. E. J., Brigham, K. S., Goldstein, M., Misra, M., & Singhal, V. (2021). Oct 7). Clinical, biochemical, and hematological characteristics of community-dwelling adolescent and young adult males with anorexia nervosa. *International Journal of Eating Disorders*. https://doi.org/10.1002/eat.23622
- Skolnick, A., Schulman, R. C., Galindo, R. J., & Mechanick, J. I. (2016). Fall). The endocrinopathies of male anorexia nervosa: Case series. AACE Clin Case Rep, 2(4), e351–e357. https://doi. org/10.4158/EP15945.CR
- Spillane, N. S., Boerner, L. M., Anderson, K. G., & Smith, G. T. (2004). Comparability of the eating disorder inventory-2 between women and men. *Assessment*, 11(1), 85–93. https://doi. org/10.1177/1073191103260623
- Støving, R. K. (2019). Mechanisms in endocrinology: Anorexia nervosa and endocrinology: A clinical update. *European Journal of Endocrinology*, 180(1), R9-r27. https://doi.org/10. 1530/eje-18-0596
- Toifl, K., Waldhauser, F., Lischka, A., & Frisch, H. (1988).
 [Anorexia nervosa in male adolescents. II. Psychoneuroendocrinologic findings]. *Klinische Pädiatrie*, 200(4), 316–320.
 (Anorexia nervosa bei männlichen Jugendlichen. Teil II: Psychoneuroendokrinologische Befunde). https://doi.org/10. 1055/s-2008-1033727
- Tomba, E., Tecuta, L., Crocetti, E., Squarcio, F., & Tomei, G. (2019). Residual eating disorder symptoms and clinical features in remitted and recovered eating disorder patients: A systematic review with meta-analysis. *International Journal of Eating Disorders*, 52(7), 759–776. https://doi.org/10.1002/eat. 23095
- van Eeden, A. E., van Hoeken, D., & Hoek, H. W. (2021). Nov 1). Incidence, prevalence and mortality of anorexia nervosa and bulimia nervosa. *Current Opinion in Psychiatry*, *34*(6), 515–524. https://doi.org/10.1097/yco.00000000000739

- Journal of Adolescent Health, 59(4), 397–400. https://doi.org/ 10.1016/j.jadohealth.2016.04.005
 Wabitsch, M., Ballauff, A., Holl, R., Blum, W. F., Heinze, E., Remschmidt, H., & Hebebrand, J. (2001). Serum leptin, gonadotropin, and testosterone concentrations in male patients with anorexia nervosa during weight gain. Journal of Clinical Endocrinology & Metabolism, 86(7), 2982–2988. https://doi.org/10.1210/jcem.86.7.7685
 Waldorf, M., Cordes, M., Vocks, S., & McCreary, D. (2014). Drive for muscularity scale--German version. PsycTESTS Dataset. https://doi.org/10.6102/zis246
 Wickham, H. (2016). ggplot2: Elegant graphics for data analysis. Springer-Verlag New York. Retrieved from https://ggplot2.
- tidyverse.org
 Wronski, M. L., Tam, F. I., Seidel, M., Mirtschink, P., Poitz, D. M., Bahnsen, K., Steinhäuser, L., Bauer, M., Roessner, V., & Ehrlich, S. (2021). Associations between pituitary-thyroid hormones and depressive symptoms in individuals with anorexia nervosa before and after weight-recovery. *Psychoneuroendocrinology*, *137*, 105630. https://doi.org/10.1016/j.psyneuen.2021.105630

Vo, M., Lau, J., & Rubinstein, M. (2016). Eating disorders in

adolescent and young adult males: Presenting characteristics.

Zhang, C. (2014). What can we learn from the history of male anorexia nervosa? *J Eat Disord*, 2(1), 138. https://doi.org/10. 1186/s40337-014-0036-9

SUPPORTING INFORMATION

Additional supporting information can be found online in the Supporting Information section at the end of this article.

How to cite this article: Schloesser, L., Lotter, L. D., Offermann, J., Borucki, K., Biemann, R., Seitz, J., Konrad, K., & Herpertz-Dahlmann, B. (2023). Sex-dependent clinical presentation, body image, and endocrine status in long-term remitted anorexia nervosa. *European Eating Disorders Review*, 1–13. https://doi.org/10.1002/erv.2997