

Results: Discovery

- Sample: Young Adult Human Connectome Project⁸, n = 143, 20 35 years.
- Focus on regularized partial correlation connectomes, which were more sensitive and reliable as compared to Pearson connectomes (see above).

Results: Application

- Sample: Early Psychosis Human Connectome Project⁹, n = 96 with schizophrenia spectrum diagnoses (non-affective: SCZ, affective: SCA), n = 55 controls (CTRL).
- All effects observed in the discovery sample replicated in the CTRL cohort.

sex, site, and motion.



First column: AUC (bars: mean & 90% percentile intervals; violins: null distribution). Second & third columns: Retest reliability (intraclass correlation and coefficient of

variance).

Percentile



nih_totalcogcomp_unadjusted



References: (1) Thiebaut de Schotten, M., & Forkel, S. J. (2022). Science | (2) Dipasquale, O. et al. (2023). Nature Neuroscience. | (4) Dworetsky, A. et al. (2021). NeuroImage. | (5) Hermosillo, R. J. M. et al. (2024). Nature Neuroscience. | (6) Markello, R. D. et al. (2022). Nature Methods. | (7) Dukart, J. et al. (2021). Human Brain Mapping | (8) Van Essen, D. C. et al. (2013). NeuroImage. | (9) Jacobs, G. R. et al. (2024). Schizophrenia Bulletin.

Acknowledgments: Leon D. Lotter: supported by BMBF and MPG, Germany. | HCP-YA: institutions supporting NIH Blueprint for Neuroscience Research (1U54MH091657); McDonnell Center. | HCP-EP: National Institute of Mental Health (U01MH109977).